

The background of the slide is a deep space image featuring two prominent purple nebulae. One is on the left, and the other is on the right, both with bright, glowing cores and diffuse, wispy edges. The space between them is filled with numerous stars of varying colors, including white, yellow, and blue, some with prominent diffraction spikes. The overall scene is dark, emphasizing the vibrant colors of the nebulae and the points of light.

High Energy Astrophysics – Status and Perspectives


Christoph Pfrommer

Heidelberg Institute for Theoretical Studies, Germany


*Annual Meeting of the German Astronomical Society,
Bochum University, 2016*

Which astrophysics can we probe at high energies?


active galactic nuclei




intergalactic space




galaxy formation



... and don't forget the UNEXPECTED!



dark matter



*structure of
space time*



*particle acceleration
magnetic amplification*

The Questions

Probing physics and cosmology with in high-energy astrophysics

- **which objects can we see?**
active galactic nuclei (blazars, radio galaxies), starburst galaxies, gamma-ray bursts, diffuse radiation, compact objects
→ astronomy: characterization, population studies
- **what underlying physics can we probe?**
most extreme physics laboratories of the cosmos:
plasma instabilities, particle acceleration, magnetic fields
→ high-energy astrophysics, plasma physics
- **what (fundamental) physics can we hope to learn?**
galaxy formation, dark matter, structure of space time
→ structure formation, cosmology, particle physics



Radiative processes induced by cosmic rays


Complementary information to cosmic rays: non-thermal emission points back to origin

hadronic processes:

- pion decay:



- photo-meson production:



- Bethe-Heitler pair production:



leptonic processes:

- inverse Compton:




- synchrotron radiation:




- bremsstrahlung:




A sketch of the non-thermal emission



The physics and cosmology of active galactic nuclei




The physics and cosmology of active galactic nuclei



Blazar: jet aligned with line-of-sight


Active galactic nuclei: paradigm and open questions




- **paradigm for γ -ray emission:**
 - synchrotron self Compton
 - external Compton
 - proton-induced cascades
 - proton synchrotron
- **open questions:**
 - energetics
 - mechanisms for jet formation and collimation
 - plasma composition (leptonic vs. hadronic, 1-zone vs. spine-layer)
 - acceleration mechanisms
- **TeV “flares”** may sign plasma instabilities in the black hole accretion disk or magnetic reconnection events in the jet



Feedback heating: M87 at radio wavelengths



$\nu = 1.4$ GHz (Owen+ 2000)




$\nu = 140$ MHz (LOFAR/de Gasperin+ 2012)

- high- ν : freshly accelerated CR electrons
low- ν : fossil CR electrons \rightarrow time-integrated AGN feedback!
- LOFAR: same picture \rightarrow puzzle of “missing fossil electrons”
- solution: electrons are fully mixed with the dense cluster gas and cooled through Coulomb interactions

The gamma-ray picture of M87

- **high state** is time variable
 → jet emission
- **low state:**
 - (1) steady flux
 - (2) γ -ray spectral index (2.2)
 = CRp index
 = CRe injection index as probed by LOFAR
 - (3) spatial extension is under investigation (?)



Rieger & Aharonian (2012)

→ **confirming this triad would be smoking gun for first γ -ray signal from a galaxy cluster!**




AGN feedback = cosmic ray heating (?)

hypothesis: low state γ -ray emission traces π^0 decay within cluster

- cosmic rays excite Alfvén waves that dissipate the energy \rightarrow heating rate

$$\mathcal{H}_{\text{CR}} = |\mathbf{v}_A \cdot \nabla P_{\text{CR}}|$$

- calibrate P_{CR} to γ -ray emission and \mathbf{v}_A to radio and X-ray emission
 \rightarrow spatial heating profile



CP (2013)


\rightarrow cosmic-ray heating matches radiative cooling (observed in X-rays) and may solve the famous “cooling flow problem” in galaxy clusters!



AGN physics and cosmology
Propagation of gamma rays
Supernova remnants

Physics
Feedback heating
Structure of space time

Probing the structure of space-time with gamma rays



Probing the structure of space-time: idea

- does quantum gravity make space-time ‘foamy’ or discrete at the Planck scale?

$$l_P = \hbar/(m_P c), \quad t_P = \hbar/(m_P c^2), \quad m_P = \sqrt{\hbar c/G}$$


- this does not happen in string theory, but in other approaches like *loop quantum gravity*
- preserving the $O(3)$ subgroup of $SO(3, 1)$, we parametrize the new dispersion rel. for photons

$$c^2 \mathbf{p}^2 = E^2 (1 + \xi E/E_{QG} + \eta E^2/E_{QG}^2 + \dots)$$

- assuming the Hamiltonian equ. of motions $\dot{x}_i = \partial H / \partial p_i$, we get

$$v \equiv \partial E / \partial p = c (1 - \xi E/E_{QG} + \dots) \Rightarrow \Delta t = \xi E/E_{QG} L/c$$

→ we can test this *energy-dependent time delay* by studying the propagation of high-energy gamma ray pulses (Amelino-Camelia+ 1998)



Quantum gravity constraints with gamma-ray bursts


- **expected time delay** for $E_{QG} \sim E_P = 10^{19}$ GeV and GeV pulse structure

$$\Delta t \approx 10 \text{ ms} \frac{E}{\text{GeV}} \frac{L}{\text{Gpc}}$$


- **idea: use pulses from gamma-ray bursts or blazar flares**
- assuming anomalous photon dispersion dominated by the linear term yields the constraint (Abdo+ 2009)

$$E_{QG} > 1.2 \times 10^{19} \text{ GeV, for } \xi = 1$$

... set mainly by the early arrival time of the 31 GeV photon!



Quantum gravity constraints with blazar flares




- no observable time delay between low and high energy photons!
- constraints on energy-dependent violation of Lorentz invariance:
 $E_{\text{QG}} > 2.1 \times 10^{18}$ GeV (90% CL limit)
- photons of all energies travel in vacuum at the same speed!

AGN physics and cosmology
Propagation of gamma rays
Supernova remnants

Extragalactic background light
Intergalactic magnetic fields?
Plasma physics

Propagation of γ rays through intergalactic space




Illustris simulation, Vogelsberger+ (2014)



High Energy Astrophysics


Observational gamma-ray cosmology

Annihilation and pair production




Observational gamma-ray cosmology

Annihilation and pair production



The *Fermi* gamma-ray horizon




Ackermann+ (2012)

- stacking of 150 significantly detected BL Lac blazars
- absorption feature moves to lower E for higher source redshifts (propagation distances) due to attenuation of gamma rays by EBL



Extragalactic background light


Unique probe of the integrated star formation rate




Dwek & Krennrich (2012)




Annihilation and pair production



Inverse Compton cascades




Inverse Compton cascades



→ each TeV point source should also be a GeV point source!


What about the cascade emission?

Every TeV source should be associated with a 1-100 GeV gamma-ray halo




What about the cascade emission?


Every TeV source should be associated with a 1-100 GeV gamma-ray halo – **not seen!**




Inverse Compton cascades



Magnetic field deflection




Magnetic field deflection




- GeV point source diluted \rightarrow weak "pair halo"
- stronger B-field implies more deflection and dilution, gamma-ray non-detection $\rightarrow B \gtrsim 10^{-16}$ G – primordial fields?

Magnetic field deflection




- **problem for unified AGN model:** no increase in comoving blazar density with redshift allowed (as seen in other AGNs) since otherwise, extragalactic GeV background would be overproduced!

What else could happen?



Plasma instabilities




→ pair plasma beam propagating through the intergalactic medium

Plasma instabilities

- **pair beam**

intergalactic medium (IGM)




- this configuration is unstable to **plasma instabilities**
- characteristic frequency and length scale of the problem:

$$\omega_p = \sqrt{\frac{4\pi e^2 n_e}{m_e}}, \quad \lambda_p = \frac{c}{\omega_p} \Big|_{\bar{\rho}(z=0)} \sim 10^8 \text{ cm}$$



Beam physics – linear growth rates

excluded for collective
 plasma phenomena



Broderick, Chang, CP (2012), also Schlickeiser+ (2012)

- consider a light beam penetrating into relatively dense plasma

- maximum growth rate

$$\Gamma \simeq 0.4 \gamma \frac{n_{\text{beam}}}{n_{\text{IGM}}} \omega_p$$

- oblique instability beats inverse Compton cooling by factor 10-100

- **assume** that instability grows at linear rate up to saturation



TeV emission from blazars – a new paradigm



absence of γ_{GeV} 's has significant implications for ...


- intergalactic magnetic field estimates
- unified picture of TeV blazars and quasars:
explains *Fermi's* γ -ray background and blazar number counts

additional IGM heating has significant implications for ...

- thermal history of the IGM: Lyman- α forest
- late time structure formation: dwarf galaxies, galaxy clusters



Extragalactic gamma-ray background




→ evolving population of hard blazars provides excellent match to latest EGRB by *Fermi* for $E \gtrsim 3$ GeV



Supernova remnants probe acceleration physics

How high-energy astrophysics impacts on cosmological structure formation

supernova remnant SN1006



X-ray: NASA/radio: NRAO/optical: NOAO




starburst galaxy M82


Supernova remnants probe acceleration physics

- **supernova remnant shocks amplify magnetic fields and accelerate CR electrons up to ~ 100 TeV** (narrow X-ray synchrotron filaments observed by *Chandra*)
- **pion bump provides evidence for CR proton acceleration** (*Fermi*/AGILE γ -ray spectra)

Fermi observations of W44:




Ackermann+ (2013)



Supernova remnants probe acceleration physics


- **supernova remnant shocks amplify magnetic fields and accelerate CR electrons up to ~ 100 TeV** (narrow X-ray synchrotron filaments observed by *Chandra*)
- **pion bump provides evidence for CR proton acceleration** (*Fermi*/AGILE γ -ray spectra)
- **shell-type SNRs show evidence for efficient shock acceleration beyond ~ 100 TeV** (HESS TeV γ -ray observations)

Fermi observations of W44:



Ackermann+ (2013)

HESS observations of shell-type SNRs:



Hinton (2009)



Physics of galaxy formation



supernova Cassiopeia A

X-ray: NASA/CXC/SAO; Optical: NASA/STScI;
Infrared: NASA/JPL-Caltech/Steward/O.Krause et al.

- galactic supernova remnants drive shock waves, accelerate electrons, amplify magnetic fields

Physics of galaxy formation




super wind in M82

NASA/JPL-Caltech/STScI/CXC/UofA

- galactic supernova remnants drive shock waves, accelerate electrons, amplify magnetic fields
- star formation and supernovae drive gas out of galaxies by galactic super winds
- critical for understanding the physics of galaxy formation → explains puzzle of low star formation efficiency in dwarf galaxies

Physics of galaxy formation



"I THINK YOU SHOULD BE MORE EXPLICIT
HERE IN STEP TWO."

A 1964 NY TIMES CARTOON

Distributed by Cullen-Expressions Ltd


- galactic supernova remnants drive shock waves, accelerate electrons, amplify magnetic fields
- star formation and supernovae drive gas out of galaxies by galactic super winds
- critical for understanding the physics of galaxy formation → explains puzzle of low star formation efficiency in dwarf galaxies

Cosmic ray-driven winds



super wind in M82

NASA/JPL-Caltech/STScI/CXG/UofA



v_z of galaxy simulation, $10^{11} M_{\odot}$

Pakmor, CP+ (2016)



- **MHD-CR simulation: diffusing cosmic rays successfully launch and energize super winds** that expel gas from the halo



Starburst galaxies

M82

Both:
D ~3 Mpc
SFR \geq SFR in MW
(in a compact region)
 $F_g \sim 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$



NGC 253

Cosmic rays and star formation



the picture: star formation → supernova remnants → proton acceleration → pion decay gamma rays induced by p-p interactions

- **dense material in starburst region**

- $\langle n \rangle \sim 250 \text{ cm}^{-3}$
- $t_{\text{pp}} \sim t_{\text{esc}}$
- approaching the calorimetric limit
- large NT bremsstrahlung and B : efficient electron emission


- **far-IR – radio correlation**

- implies universal conversion: star form. → CR → synchrotron
- now: far-IR – gamma-ray correlation




Far infra-red – gamma-ray correlation

Universal conversion: star formation \rightarrow cosmic rays \rightarrow gamma rays



Far infra-red – gamma-ray correlation

Universal conversion: star formation \rightarrow cosmic rays \rightarrow gamma rays



Conclusions

- the non-thermal universe uncovered by high-energy radiation provides **new probes of fundamental physics and cosmology**
- **radio and X-ray astronomy have provided impressive discoveries** of new phenomena
- **now the age of cosmic-ray and γ -ray astronomy has begun** and neutrino (and gravitational wave) astronomy are about to open up

→ **non-thermal multi-messenger analyses:**

“The only true voyage of discovery would be not to visit new landscapes but to possess other eyes and to behold the universe through the eyes of another, of a hundred others.”

Marcel Proust



AGN physics and cosmology
Propagation of gamma rays
Supernova remnants

Particle acceleration
Galaxy formation
Starburst galaxies

CRAGSMAN: The Impact of Cosmic RAYs on Galaxy and CluSTER ForMAtion



This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No CRAGSMAN-646955).