



*30,000 foot view of blazar heating*

Christoph Pfrommer<sup>1</sup>

with


Avery E. Broderick, Phil Chang, Ewald Puchwein,  
Astrid Lamberts, Mohamad Shalaby, Volker Springel

<sup>1</sup>Heidelberg Institute for Theoretical Studies, Germany

Feedback over 44 orders of magnitude, Perimeter Institute – 2016


# Motivation

A new link between high-energy astrophysics and cosmological structure formation




- **Introduction to Blazars**
  - active galactic nuclei (AGN)
  - propagating gamma rays
  - plasma physics
- **Cosmological Consequences**
  - unifying blazars with AGN
  - gamma-ray background
  - thermal history of the Universe
  - Lyman- $\alpha$  forest
  - formation of dwarf galaxies

# Active galactic nucleus (AGN)



- **AGN: compact region at the center of a galaxy**, which dominates the luminosity of its electromagnetic spectrum
- AGN emission is most likely caused by **mass accretion onto a supermassive black hole** and can also launch **relativistic jets**


# Active galactic nucleus at a cosmological distance




Quasar 3C175 at  $z \simeq 0.8$ :  
jet extends  $10^6$  light years across

- **AGN: compact region at the center of a galaxy**, which dominates the luminosity of its electromagnetic spectrum
- AGN emission is most likely caused by **mass accretion onto a supermassive black hole** and can also launch **relativistic jets**
- AGNs are among the most luminous sources in the universe  
→ **discovery of distant objects**

# Unified model of active galactic nuclei



# Unified model of active galactic nuclei



Blazar: jet aligned with line-of-sight

Blazars  
Gamma-ray sky  
Structure formation

Active galactic nuclei  
Propagating  $\gamma$  rays  
Plasma instabilities

# TeV gamma-ray observations



MAGIC



VERITAS




H.E.S.S.



# The TeV gamma-ray sky


There are several classes of TeV sources:

- Galactic - pulsars, BH binaries, supernova remnants
- Extragalactic - **mostly** blazars, two starburst galaxies







# Annihilation and pair production




# Annihilation and pair production



# Inverse Compton cascades




# Inverse Compton cascades



→ each TeV point source should also be a GeV point source!


# What about the cascade emission?

Every TeV source should be associated with a 1-100 GeV gamma-ray halo




# What about the cascade emission?


Every TeV source should be associated with a 1-100 GeV gamma-ray halo – **not seen!**



# Inverse Compton cascades




# Extragalactic magnetic fields?






# Extragalactic magnetic fields?




- GeV point source diluted  $\rightarrow$  weak "pair halo"
- stronger B-field implies more deflection and dilution, gamma-ray non-detection  $\rightarrow B \gtrsim 10^{-16}$  G – primordial fields?

# Extragalactic magnetic fields?




- **problem for unified AGN model:** no increase in comoving blazar density with redshift allowed (as seen in other AGNs) since otherwise, extragalactic GeV background would be overproduced!

# What else could happen?



# Plasma instabilities




→ pair plasma beam propagating through the intergalactic medium

# Plasma instabilities

- **pair beam**

- **intergalactic medium (IGM)**



- this configuration is unstable to **plasma instabilities**
- characteristic frequency and length scale of the problem:


$$\omega_p = \sqrt{\frac{4\pi e^2 n_e}{m_e}}, \quad \lambda_p = \frac{c}{\omega_p} \Big|_{\bar{\rho}(z=0)} \sim 10^8 \text{ cm}$$



# Two-stream instability

consider wave-like perturbation in background plasma along the beam direction (Langmuir wave):


- initially homogeneous beam- $e^-$ :  
attractive (repulsive) force by potential maxima (minima)
- $e^-$  attain lowest velocity in potential minima  $\rightarrow$  bunching up
- $e^+$  attain lowest velocity in potential maxima  $\rightarrow$  bunching up




# Two-stream instability

consider wave-like perturbation in background plasma along the beam direction (Langmuir wave):

- beam- $e^+/e^-$  couple in phase with the background perturbation: enhances background potential
- stronger forces on beam- $e^+/e^- \rightarrow$  positive feedback
- exponential wave-growth  $\rightarrow$  instability



# Two-stream instability: momentum transfer





- particles with  $v \gtrsim v_{phase}$ :  
pair momentum  $\rightarrow$  plasma waves  $\rightarrow$  growing modes: instability
- particles with  $v \lesssim v_{phase}$ :  
plasma wave momentum  $\rightarrow$  pairs  $\rightarrow$  Landau damping





# Oblique instability


- $\mathbf{k}$  oblique to  $\mathbf{v}_{\text{beam}}$ : real world perturbations don't choose "easy" alignment =  $\sum$  all orientations
- **oblique grows faster than two-stream**:  $E$ -fields can easier deflect ultra-relativistic particles than change their parallel velocities  
(Nakar, Bret & Milosavljevic 2011)



Bret (2009), Bret+ (2010)

# Beam physics – growth rates

excluded for collective  
plasma phenomena



Broderick, Chang, C.P. (2012), also Schlickeiser+ (2012)

- consider a light beam penetrating into relatively dense plasma

- maximum growth rate


$$\Gamma \simeq 0.4 \gamma \frac{n_{\text{beam}}}{n_{\text{IGM}}} \omega_p$$

- oblique instability beats inverse Compton cooling by factor 10-100

- **assume** that instability grows at *linear* rate up to saturation



# TeV emission from blazars – a new paradigm




absence of  $\gamma_{\text{GeV}}$ 's has significant implications for ...


- intergalactic magnetic field estimates
- unified picture of TeV blazars and quasars



# Implications for intergalactic magnetic fields



- competition of rates:  
 $\Gamma_{\text{IC}}$  vs.  $\Gamma_{\text{oblique}}$
- fraction of the pair energy lost to inverse-Compton on the CMB:  
 $f_{\text{IC}} = \Gamma_{\text{IC}} / (\Gamma_{\text{IC}} + \Gamma_{\text{oblique}})$
- **plasma instability dominates for more luminous blazars**



Broderick, Chang, C.P. (2012)



## Conclusions on $B$ -field constraints from blazar spectra


- it is thought that TeV blazar spectra might constrain IGM  $B$ -fields
- this assumes that cooling mechanism is IC off the CMB + deflection from magnetic fields
- beam instabilities allow high-energy  $e^+ / e^-$  pairs to self scatter and/or lose energy
- isotropizes the beam – no need for  $B$ -field
- $\lesssim 1\text{--}10\%$  of beam energy to IC CMB photons

→ **TeV blazar spectra are not suitable to measure IGM  $B$ -fields**  
(if plasma instabilities saturate close to linear rate)!

Broderick, Chang, C.P. (2012), Schlickeiser, Krakau, Supsar (2013), Chang+ (2014)



# TeV blazar luminosity density: today




Broderick, Chang, C.P. (2012)

- collect luminosity of all 23 TeV blazars with good spectral measurements
- account for the selection effects (sky coverage, duty cycle, galactic occultation, TeV flux limit)
- TeV blazar luminosity density is a scaled version ( $\eta_B \sim 0.2\%$ ) of that of quasars!



# Unified TeV blazar-quasar model



Broderick, Chang, C.P. (2012)

Quasars and TeV blazars are:

- regulated by the same mechanism
- contemporaneous elements of a single AGN population: TeV-blazar activity does not lag quasar activity

→ **assume that they trace each other for all redshifts!**




# How many TeV blazars are there?



→ use all-sky survey of  
the GeV gamma-ray sky:  
*Fermi* gamma-ray space  
telescope




# How many TeV blazars are there?



Hopkins+ (2007)




# How many TeV blazars are there?



Hopkins+ (2007)




# How many TeV blazars are there?



Hopkins+ (2007)




# Redshift distribution of *Fermi* hard $\gamma$ -ray blazars



→ evolving (increasing) blazar population consistent with observed declining evolution (*Fermi* flux limit)!




# $\log \mathcal{N} - \log S$ distribution of *Fermi* hard $\gamma$ -ray blazars



→ predicted and observed flux distributions of hard *Fermi* blazars between 10 GeV and 500 GeV are indistinguishable!




# How many TeV blazars are there?



Hopkins+ (2007)




# TeV photon absorption by pair production



intrinsic and **observed** SEDs  
of blazars at  $z = 1$

→  $\gamma$ -ray attenuation by an-  
nihilation and pair producing  
on the EBL



inferred spectral index  $\Gamma_F$  for  
the spectra in the top panel;  
overlay of *Fermi* data on  
BL Lacs and non-BL Lacs  
(mostly FSRQs)

Broderick, C.P.+ (2013)



# Extragalactic gamma-ray background

- intrinsic spectrum for a TeV blazar:

$$\frac{dN}{dE} = f \hat{F}_E = f \left[ \left( \frac{E}{E_b} \right)^{\Gamma_l} + \left( \frac{E}{E_b} \right)^{\Gamma_h} \right]^{-1},$$

$E_b = 1$  TeV is break energy,  $\Gamma_h = 3$  is high-energy spectral index,  
 $\Gamma_l$  related to  $\Gamma_F$ , which is drawn from observed distribution

- extragalactic gamma-ray background (EGRB):

$$E^2 \frac{dN}{dE}(E, z) = \frac{1}{4\pi} \int_0^2 d\Gamma_l \int_z^\infty dV(z') \frac{\eta_B \tilde{\Lambda}_Q(z') \hat{F}_{E'}}{4\pi D_L^2} e^{-\tau_E(E', z')},$$

$E' = E(1 + z')$  is gamma-ray energy at *emission*,


$\tilde{\Lambda}_Q$  is physical quasar luminosity density,

$\eta_B \sim 0.2\%$  is blazar fraction,  $\tau$  is optical depth






# Extragalactic gamma-ray background



→ evolving population of hard blazars provides excellent match to latest EGRB by *Fermi* for  $E \gtrsim 3$  GeV




# Extragalactic gamma-ray background



→ the signal at 10 (100) GeV is dominated by redshifts  $z \sim 1.2$  ( $z \sim 0.6$ )



# TeV emission from blazars – a new paradigm



absence of  $\gamma_{\text{GeV}}$ 's has significant implications for ...

- intergalactic magnetic field estimates
- unified picture of TeV blazars and quasars:  
explains *Fermi's*  $\gamma$ -ray background and blazar number counts

additional IGM heating has significant implications for ...

- thermal history of the IGM: Lyman- $\alpha$  forest
- late-time formation of dwarf galaxies



# Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars
- $T_{\text{IGM}} \sim 10^4$  K (1 eV) at mean density ( $z \sim 2$ )

$$\varepsilon_{\text{th}} = \frac{kT}{m_p c^2} \sim 10^{-9}$$

- radiative energy ratio emitted by BHs in the Universe (Fukugita & Peebles 2004)

$$\varepsilon_{\text{rad}} = \eta \Omega_{\text{bh}} \sim 0.1 \times 10^{-4} \sim 10^{-5}$$

- fraction of the energy energetic enough to ionize H I is  $\sim 0.1$ :


$$\varepsilon_{\text{UV}} \sim 0.1 \varepsilon_{\text{rad}} \sim 10^{-6} \quad \rightarrow \quad kT \sim \text{keV}$$

- photoheating efficiency  $\eta_{\text{ph}} \sim 10^{-3} \quad \rightarrow \quad kT \sim \eta_{\text{ph}} \varepsilon_{\text{UV}} m_p c^2 \sim \text{eV}$   
(limited by the abundance of H I/He II due to the small recombination rate)

- blazar heating efficiency  $\eta_{\text{bh}} \sim 10^{-3} \quad \rightarrow \quad kT \sim \eta_{\text{bh}} \varepsilon_{\text{rad}} m_p c^2 \sim 10 \text{ eV}$   
(limited by the total power of TeV sources)



# Thermal history of the IGM




C.P., Chang, Broderick (2012)

→ increased temperature at **mean** density!




# Evolution of the temperature-density relation

no blazar heating



with blazar heating




Chang, Broderick, C.P. (2012)

- blazars and extragalactic background light are uniform:
  - blazar heating rate independent of density
  - makes low density regions *hot*
  - causes inverted temperature-density relation,  $T \propto 1/\delta$




# Blazars cause hot voids

no blazar heating



with blazar heating




Chang, Broderick, C.P. (2012)

- blazars completely change the thermal history of the diffuse IGM and late-time structure formation




# Cosmological hydrodynamical simulations

- include predicted volumetric heating rate in cosmological hydrodynamical simulations
- study:
  - thermal properties of intergalactic medium
  - Lyman- $\alpha$  forest






# Temperature-density relation




Puchwein, C.P., Springel, Broderick, Chang (2012)




# The Lyman- $\alpha$ forest



# The observed Lyman- $\alpha$ forest





# The simulated Ly- $\alpha$ forest



Puchwein, C.P.+ (2012)



# Optical depths and temperatures




Puchwein, C.P.+ (2012)


Redshift evolutions of effective optical depth and IGM temperature match data only with additional heating, e.g., provided by blazars!



# Ly- $\alpha$ flux PDFs and power spectra



Puchwein, C.P.+ (2012)




# Lyman- $\alpha$ forest in a blazar heated Universe

improvement in modelling the Lyman- $\alpha$  forest is a direct consequence of the peculiar properties of blazar heating:

- **heating rate independent of IGM density**  $\rightarrow$  naturally produces the inverted  $T-\rho$  relation that Lyman- $\alpha$  forest data demand
- **recent and continuous nature of the heating** is needed to match the redshift evolutions of all Lyman- $\alpha$  forest statistics
- **magnitude of the heating rate required by Lyman- $\alpha$  forest data**  $\sim$  the total energy output of TeV blazars (or equivalently  $\sim 0.2\%$  of that of quasars)



# “Missing satellite” problem in the Milky Way



Substructures in cold DM simulations much more numerous than observed number of Milky Way satellites!





# Dwarf galaxy formation


- thermal pressure opposes gravitational collapse on small scales
- characteristic length/mass scale below which objects do not form
- hotter intergalactic medium  $\rightarrow$  higher thermal pressure  
 $\rightarrow$  higher Jeans mass:

$$M_J \propto \frac{c_s^3}{\rho^{1/2}} \propto \left( \frac{T_{\text{IGM}}^3}{\rho} \right)^{1/2} \rightarrow \frac{M_{J,\text{blazar}}}{M_{J,\text{photo}}} \approx \left( \frac{T_{\text{blazar}}}{T_{\text{photo}}} \right)^{3/2} \gtrsim 30$$

$\rightarrow$  blazar heating increases  $M_J$  by 30 over pure photoheating!

- complications:  
non-linear collapse,  
delayed pressure response in expanding universe  $\rightarrow$  concept of “filtering mass”

# Dwarf galaxy formation suppressed




C.P., Chang, Broderick (2012)


- blazar heating suppresses the formation of late-forming dwarfs within existing dark matter halos of masses  $< 10^{11} M_\odot$   
 → introduces new time and mass scale to galaxy formation!



# When do dwarfs form?




Dolphin+ (2005)



isochrone fitting for different metallicities → star formation histories



# When do dwarfs form?




Dolphin+ (2005)




# Milky Way satellites: formation history and abundance

satellite formation time



satellite luminosity function



Maccio+ (2010)

- blazar heating suppresses late satellite formation, may reconcile low observed dwarf abundances with CDM simulations



# Conclusions on blazar heating

**Blazar heating:** TeV photons are attenuated by EBL; their kinetic energy  $\rightarrow$  heating of the IGM; it is *not* cascaded to GeV energies

- **explains puzzles in gamma-ray astrophysics:**
  - lack of GeV bumps in blazar spectra without IGM  $B$ -fields
  - *unified TeV blazar-quasar model* explains Fermi source counts and extragalactic gamma-ray background
- **novel mechanism; dramatically alters thermal history of the IGM:**
  - uniform and  $z$ -dependent preheating
  - quantitative self-consistent picture of high- $z$  Lyman- $\alpha$  forest
- **significantly modifies late-time structure formation:**
  - suppresses late dwarf formation
  - void phenomenon, “missing satellites” (?)



# Literature for the talk

- Broderick, Chang, Pfrommer, *The cosmological impact of luminous TeV blazars I: implications of plasma instabilities for the intergalactic magnetic field and extragalactic gamma-ray background*, ApJ, 752, 22, 2012.
- Chang, Broderick, Pfrommer, *The cosmological impact of luminous TeV blazars II: rewriting the thermal history of the intergalactic medium*, ApJ, 752, 23, 2012.
- Pfrommer, Chang, Broderick, *The cosmological impact of luminous TeV blazars III: implications for galaxy clusters and the formation of dwarf galaxies*, ApJ, 752, 24, 2012.
- Puchwein, Pfrommer, Springel, Broderick, Chang, *The Lyman- $\alpha$  forest in a blazar-heated Universe*, MNRAS, 423, 149, 2012.
- Broderick, Pfrommer, Chang, Puchwein, *Implications of plasma beam instabilities for the statistics of the Fermi hard gamma-ray blazars and the origin of the extragalactic gamma-ray background*, ApJ, 790, 137, 2014.
- Chang, Broderick, Pfrommer, Puchwein, Lamberts, Shalaby, *The effect of nonlinear Landau damping on ultrarelativistic beam plasma instabilities*, ApJ, 2014, 797, 110.
- Lamberts, Chang, Pfrommer, Puchwein, Broderick, Shalaby, *Patchy blazar heating: diversifying the thermal history of the intergalactic medium*, 2015, submitted, arXiv:1502.07980.



# Additional slides






# Empirical model for star formation histories (1)

Lu, Mo, Lu, Katz, et al. (2013):  
constructing merger tree-based  
model of galaxy formation that  
matches

- observed stellar mass function (different  $z$ )
- luminosity function of local cluster galaxies


→ star formation histories of  
dark matter halos (different  $z$ )



Lu+ (2013)



## Empirical model for star formation histories (2)





Lu+ (2013)

→ strong quenching of star formation efficiency for  $z \lesssim 2$  in low-mass halos ( $M < 10^{11} h^{-1} M_{\odot}$ ) → **blazar heating?**



# Galactic H I-mass function

Mo+ (2005)



- H I-mass function is too flat (i.e., gas version of missing dwarf problem!)
- photoheating and SN feedback too inefficient
- IGM entropy floor of  $K \sim 15 \text{ keV cm}^2$  at  $z \sim 2 - 3$  successful!

